


Status of the Claims:

1. Canceled.
2. Canceled.
3. Canceled.
4. Canceled.
5. Canceled.
6. (Original) An ophthalmic lens comprising a compound of the formula:





wherein X is sulfur or oxygen;

R', R'' are the same or different and are each independently hydrogen, hydroxy, halogen, nitro, cyano, allyl, linear or branched (C₁-C₂₀)alkyl, (C₃-C₂₀)cycloalkyl, (C₁-C₂₀)alkoxy, (C₁-C₂₀)alkylacetylenyl, phenylacetylenyl, (C₁-C₂₀)alkenyl, phenylvinyl, halo(C₁-C₂₀)alkyl, halo(C₃-C₂₀)cycloalkyl, halo(C₁-C₂₀)alkoxy, aryl, aryloxy or heteroaryl optionally substituted with (C₁-C₆)alkyl or (C₁-C₆)alkoxy; arylalkyl or heteroarylalkyl; nitrogen-containing heterocyclic ring having 5 or 6 atoms optionally substituted (C₁-C₆)alkyl or (C₁-C₆)alkoxy, -N(R₁)R₂, CON(R₁)R₂, wherein R₁ and R₂ may be the same or different and are each independently hydrogen, (C₁-C₂₀)alkyl, (C₃-C₂₀)cycloalkyl, and optionally substituted phenyl; -OCOR, -COOR or -COR, wherein R represents hydrogen,

(C₁-C₂₀)alkyl, (C₃-C₂₀)cycloalkyl, or aryl or heteroaryl optionally substituted with (C₁-C₆)alkyl or (C₁-C₆)alkoxy;

A', A'' may be same or different and are each independently:



linear or branched (C₁-C₁₂)alkyl, (C₃-C₁₂)cycloalkyl, aryl(C₁-C₆)alkyl or heteroaryl(C₁-C₆)alkyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl, (C₁-C₁₂)alkoxy, halo(C₁-C₁₂)alkyl, (C₁-C₁₂)haloalkoxy, (C₁-C₁₂)alkylthio;
optionally substituted aryl groups;
optionally substituted heteroaryl groups;
a group of the following formulae:



or

wherein B is hydrogen, (C₁-C₁₂)alkyl or optionally substituted aryl;
unsubstituted or mono-substituted pyrazolyl, pyridyl, imidazolyl, pyrazolinyl, imidazolinyl, or acridinyl, each of the said substituents selected from the group consisting of (C₁-C₆)alkyl, (C₁-C₆)alkoxy, fluoro, chloro, and phenyl.

(f) a group of the following formulae:



wherein C, D may be the same or different and are each independently carbon, oxygen, (C₁-C₁₂)alkyl nitrogen, or (C₁-C₁₂)acyl nitrogen;
R₃ and R₄ are each hydrogen or (C₁-C₁₂)alkyl; and

wherein the phenyl moiety is optionally substituted with (C₁-C₁₂)alkyl, (C₁-C₁₂)alkoxy, (C₂-C₁₂)acyl, fluoro, or chloro ;

n is an integer from 1 to 8; and

m is an integer from 0 to 3.

7. (Amended) The ophthalmic lens of claim 1 ~~6~~, wherein:

X is sulfur or oxygen;

R', R'' are the same or different and are each independently hydrogen, nitro, cyano, allyl, fluoro, chloro, bromo, trifluoromethyl, trichloromethyl, pyrrolidino, piperidino, morpholino, phenyl, benzyl; linear or branched (C₁-C₆)alkyl, (C₁-C₆)alkoxy, or -OCOR or -COOR wherein R is hydrogen, (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl;

A', A'' are the same or different and are each independently:

linear or branched (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, aryl(C₁-C₄)alkyl or heteroaryl(C₁-C₄)alkyl, (C₁-C₆)alkoxy(C₁-C₆)alkyl;

unsubstituted, mono-, di-substituted aryl selected from phenyl or naphthyl;; unsubstituted or mono-substituted heteroaryl groups that are furyl, thienyl, pyreryl, indolyl, benzofuryl, benzothienyl, pyridyl, dibenzofuryl, dibenzothienyl, or carbazolyl the substituents being

nitro, amino, cyano, hydroxy, epoxy, hydroxyethoxy, methoxyethoxy, hydroxyethoxyethoxy, methoxyethoxyethoxy, fluoro, chloro, bromo, iodo, vinyl, allyl, trifluoromethyl, phenyl, (C₁-C₆)alkyl, (C₁-C₆)alkoxy, cyclo(C₃-C₆)alkyl, cyclo(C₁-C₆)alkoxy, (C₁-C₆)alkylamino, di(C₁-C₆)alkylamino, diarylamino, phenylacetylenyl, or phenylvinyl;

N(C₁-C₆)alkylpiperazino, N-aryl-piperizino, aziridino, indolino, pyrrolidino, pyrrolino, piperidino, (C₁-C₄)alkylpiperidino, di(C₁-C₄)alkylpiperidino, 4-piperidinopiperidino, morpholino, 2,6-di(C₁-C₄)alkylmorpholino, thiomorpholino, thioazolidino, tetrahydroquinolino, or pyreryl;

$N(R_1)R_2$, $CON(R_1)R_2$, wherein R_1 and R_2 are the same or different and are each independently hydrogen, (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl, phenyl or $-COR$, $-OCOR$ or $-COOR$ wherein R is hydrogen, (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl, or phenyl;

n is an integer from 1 to 6; and

m is an integer from 0 to 2.

8. (Amended) The ophthalmic lens of claim 4, wherein:

X is sulfur;

R' , R'' are the same or different and are each independently hydrogen, nitro, cyano, fluoro, chloro, bromo, pyrrolidino, piperidino, morpholino, phenyl, benzyl, (C_1-C_4) alkyl, or (C_1-C_4) alkoxy;

A' , A'' are the same or different and are each independently:

a linear or branched (C_1-C_4) alkyl, (C_3-C_6) cycloalkyl;

unsubstituted, mono-, or di-substituted phenyl the substituents being one or more of nitro, amino, acyl, cyano, methoxy, ethoxy, methoxyethoxy, fluoro, chloro, vinyl, allyl, methoxycarbonyl, ethoxycarbonyl, (C_1-C_4) alkyl, di (C_1-C_4) alkylamino, piperazino, piperidino, arylpiperidino, morpholino, pyrrolidino, aziridino, acryloxy, methacryloxy, phenylacetylenyl, phenylvinyl;

unsubstituted or mono-substituted furyl, thienyl, or pyrrol substituted with a substituent that is (C_1-C_4) alkyl or phenyl;

n is an integer from 1 to 4, and

m is, independently, integer from 0 to 2.

9. (Original) An ophthalmic lens comprising a compound selected from the group consisting of:

5,5'-Bis[3-(p-methoxyphenyl)-[3H]-naphtho[2,1-b]pyran-3-yl]-2,2'-bithiophene;

5,5'-Bis[3-(p-methylphenyl)-[3H]-naphtho[2,1-b]pyran-3-yl]-2,2'-bithiophene;

5,5'-Bis[3-(p-fluorophenyl)-[3H]-naphtho[2,1-b]pyran-3-yl]-2,2'-bithiophene;

5,5'-Bis[3-(o-fluorophenyl)-[3H]-naphtho[2,1-b]pyran-3-yl]-2,2'-bithiophene;

5,5'''-Bis[3-(p-methoxyphenyl)-[3H]-naphtho[2,1-b]pyran-3-yl]-[2,2',5',2'',5'',2''']-
quaterthiophene;
5,5'''-Bis[3-(naphthalene-2-yl)-[3H]-naphtho[2,1-b]pyran-3-yl]-[2,2',5',2'',5'',2''']-
quaterthiophene; or
5,5'''-Bis[3-(o-fluorophenyl)-[3H]-naphtho[2,1-b]pyran-3-yl]-[2,2',5',2'',5'',2''']-
quaterthiophene.

10. (Original) The ophthalmic lens of claim 6, 7, 8, or 9, further comprising a photochromic compound selected from the group consisting of naphthopyran, spirooxazine, one or more nonphotochromic dyes, or a combinations thereof.


11. (Original) A method for making a naphthopyran compound comprising the step of
coupling a propargyl alcohol with a naphthol in the presence of a catalytic amount of an acid, a solvent and 3Å molecular sieves to generate the naphthopyran compound.

REMARKS

Applicant submits that the claims, as amended, are in condition for allowance.
Entry of the amendments, and allowance of the claims are requested.

Date: October 28, 2003
Johnson & Johnson
Law Department
One Johnson & Johnson Plaza
New Brunswick, New Jersey 08933

Respectfully submitted,


Lois A. Gianneschi
Reg. No. 35,519
Attorney for Applicant
(732) 524-6351